

# Dear Parents



## Considerations for Youth Sports

As some communities in the United States begin to start youth sports activities again, the Centers for Disease Control and Prevention (CDC) offers the following considerations for ways in which youth sports organizations can protect players, families, and communities and slow the spread of the Coronavirus Disease 2019 (COVID-19). Administrators of youth sports organizations can consult with [state and local health officials](#) to determine if and how to put into place these considerations. Each community may need to make adjustments to meet its unique needs and circumstances. Implementation should be guided by what is practical, acceptable, and tailored to the needs of each community. These considerations are meant to supplement – **not replace** – any state, local, territorial, or tribal health and safety laws, rules, and regulations with which youth sports organizations must comply.

## Guiding Principles to Keep in Mind

There are a number of actions youth sports organizations can take to help lower the risk of COVID-19 exposure and reduce the spread during competition and practice. The more people a child or coach interacts with, the closer the physical interaction, the more sharing of equipment there is by multiple players, and the longer that interaction, the higher the risk of COVID-19 spread. Therefore, risk of COVID-19 spread can be different, depending on the type of activity. The risk of COVID-19 spread increases in youth sports settings as follows:

- **Lowest Risk:** Performing skill-building drills or conditioning at home, alone or with family members.
- **Increasing Risk:** Team-based practice.
- **More Risk:** Within-team competition.
- **Even More Risk:** Full competition between teams from the same local geographic area.
- **Highest Risk:** Full competition between teams from different geographic areas.

# Dear Parents

If organizations are not able to keep in place safety measures during competition (for example, maintaining social distancing by keeping children six feet apart at all times), they may consider dropping down a level and limiting participation to within-team competition only (for example, scrimmages between members of the same team) or team-based practices only. Similarly, if organizations are unable to put in place safety measures during team-based activities, they may choose individual or at-home activities, especially if any members of the team are at [high-risk for severe illness](#).

## Assessing Risk

The way sports are played, and the way equipment is shared can influence the spread of COVID-19 among players. When you are assessing the risk of spread in your sport, consider:

- **Physical closeness of players, and the length of time that players are close to each other or to staff.** Sports that require frequent closeness between players may make it more difficult to maintain social distancing, compared to sports where players are not close to each other. For close-contact sports (e.g., wrestling, basketball), play may be modified to safely increase distance between players.
  - For example, players and coaches can:
    - focus on individual skill building versus competition;
    - limit the time players spend close to others by playing full contact only in game-time situations;
    - decrease the number of competitions during a season.

Coaches can also modify practices so players work on individual skills, rather than on competition. Coaches may also put players into small groups (cohorts) that remain together and work through stations, rather than switching groups or mixing groups.

- **Amount of necessary touching of shared equipment and gear (e.g., protective gear, balls, bats, racquets, mats, or water bottles).** It is also possible that a person can get COVID-19 by touching a surface or object that has the virus on it, and then touching their own mouth, nose, or eyes. Minimize equipment sharing, and clean and disinfect shared equipment between use by different people to reduce the risk of COVID-19 spread.
- **Ability to engage in social distancing while not actively engaged in play (e.g., during practice, on the sideline, or in the dugout).** During times when players are not actively participating in practice or competition, attention should be given to maintaining [social distancing](#) by increasing space between players on the sideline, dugout, or bench. Additionally, coaches can encourage athletes to use downtime for individual skill-building work or cardiovascular conditioning, rather than staying clustered together.
- **Age of the player.** Older youth might be better able to follow directions for social distancing and take other protective actions like not sharing water bottles. If feasible, a coach, parent, or other caregiver can assist with making sure that athletes maintain proper social distancing. For younger athletes, youth sports programs may ask parents or other household members to monitor their children and make sure that they follow social distancing and take other protective actions (e.g., younger children could sit with parents or caregivers, instead of in a dugout or group area).
- **Players at higher risk of developing serious disease.** Parents and coaches should assess level of risk based on individual players on the team who may be at [higher risk for severe illness](#), such as children who may have asthma, diabetes, or other health problems.

# Dear Parents

- **Size of the team.** Sports with a large number of players on a team may increase the likelihood of spread, compared to sports with fewer team members. Consider decreasing team sizes, as feasible.
- **Nonessential visitors, spectators, volunteers.** Limit any nonessential visitors, spectators, volunteers, and activities involving external groups or organizations.
- **Travel outside of the local community.** [Traveling](#) outside of the local community may increase the chances of exposing players, coaches, and fans to COVID-19, or unknowingly spreading it to others. This is the case particularly if a team from an area with high levels of COVID-19 competes with a team from an area with low levels of the virus. Youth sports teams should consider competing only against teams in their local area (e.g., neighborhood, town, or community).

## Promoting Behaviors that Reduce Spread

Youth sports organizations may consider implementing several strategies to encourage behaviors that reduce the spread of COVID-19.

- **Staying Home when Appropriate**
  - Educate staff and player families about when they should stay home and when they can return to activity
    - Actively encourage sick staff, families, and players to stay home. Develop policies that encourage sick employees to stay at home without fear of reprisal, and ensure employees aware of these policies.
    - [Individuals, including coaches, players, and families, should stay home](#) if they have tested positive for or are showing COVID-19 [symptoms](#).
    - Individuals, including coaches, players, and families, who have recently had a [close contact](#) with a person with COVID-19 should also [stay home and monitor their health](#).
    - CDC's criteria can help inform return to work/school policies:
    - [If they have been sick with COVID-19](#)
    - [If they have recently had a close contact with a person with COVID-19](#)
- **Hand Hygiene and Respiratory Etiquette**
  - Teach and reinforce [handwashing](#) with soap and water for at least 20 seconds
    - If soap and water are not readily available, hand sanitizer that contains at least 60% alcohol can be used (for staff and older children who can safely use hand sanitizer).
  - Do not allow spitting and encourage everyone to cover their coughs and sneezes with a tissue or use the inside of their elbow. Used tissues should be thrown in the trash and hands washed immediately with soap and water for at least 20 seconds.
    - If soap and water are not readily available, hand sanitizer that contains at least 60% alcohol can be used.
- **Masks**
  - Teach and reinforce the use of [masks](#). Masks are not intended to protect the wearer, but rather to reduce the risk of spreading COVID-19 from the person wearing the mask (who may not have any symptoms of disease). Masks may be challenging for players (especially younger players) to wear while playing sports. Masks should be worn by coaches, youth sports staff, officials, parents, and spectators as much as possible.

# Dear Parents

- Wearing masks is most important when physical distancing is difficult.
- People wearing masks should be reminded to not touch the mask and to [wash their hands](#) frequently. Information should be provided to all participants on the [proper use, removal, and washing of masks](#).
  - Note: Masks should **not** be placed on:
    - Babies and children younger than 2 years old;
    - Anyone who has trouble breathing or is unconscious;
    - Anyone who is incapacitated or otherwise unable to remove the mask without assistance.
- **Adequate Supplies**
  - If hand washing facilities are available, support healthy hygiene by providing supplies including soap, paper towels, tissues, and no-touch/foot pedal trash cans. If hand washing facilities are not available, provide hand sanitizer with at least 60% alcohol (for coaches, staff and older players who can safely use hand sanitizer).
- **Signs and Messages**
  - Post signs in highly visible locations (e.g., at entrances and exits, and in restrooms) that describe how to [stop the spread](#) of germs such as by [properly washing hands](#) and [properly wearing a mask](#).
  - Broadcast [regular announcements on public announcement \(PA\) system](#)
  - Include COVID-19 prevention messages (for example, [videos](#)) about behaviors that prevent spread of COVID-19 when communicating with staff, volunteers, officials, and families. This could include links, videos, and prevention messages in emails, on organization websites, and through the team and league's [social media accounts](#).
  - Find freely available CDC print and digital resources on [CDC's communication resources](#) main page.

## Maintaining Healthy Environments

Youth sports organizations may consider implementing several strategies to maintain healthy environments.

- **Cleaning and Disinfection**
  - [Clean and disinfect](#) frequently touched surfaces on the field, court, or play surface (e.g., drinking fountains) at least daily, or between uses as much as possible. Use of shared objects and equipment (e.g., balls, bats, gymnastics equipment) should be limited, or cleaned between use by each individual if possible.
  - Develop a schedule for increased, routine cleaning and disinfection.
  - Ensure [safe and correct use](#) and storage of disinfectants, including storing products securely away from children. Use products that meet [EPA disinfection criteria](#).
  - Identify an adult staff member or volunteer to ensure proper cleaning and disinfection of objects and equipment, particularly for any shared equipment or frequently touched surfaces.
  - Cleaning products should not be used near children, and staff should ensure that there is adequate ventilation when using these products to prevent children or themselves from inhaling toxic fumes.
  - Use gloves when removing garbage bags or handling and disposing of trash. [Wash hands after removing gloves](#).

# Dear Parents

- **Shared Objects**
  - Discourage sharing of items that are difficult to clean, sanitize, or disinfect. Do not let players share towels, clothing, or other items they use to wipe their faces or hands.
  - Make sure there are adequate supplies of shared items to minimize sharing of equipment to the extent possible (e.g., protective gear, balls, bats, water bottles); otherwise, limit use of supplies and equipment to one group of players at a time and clean and disinfect between use.
    - Keep each player's belongings separated from others' and in individually labeled containers, bags, or areas.
    - If food is offered at any event, have pre-packaged boxes or bags for each attendee instead of a buffet or family-style meal. Avoid sharing food and utensils. Offer hand sanitizer or encourage hand washing.
- **Ventilation**
  - If playing inside, ensure ventilation systems or fans operate properly. Increase circulation of outdoor air as much as possible, for example by opening windows and doors. Do not open windows and doors if doing so poses a safety or health risk (e.g., risk of falling or triggering asthma symptoms) to players or others using the facility.
- **Water Systems ( Individual water Bottles)**
  - To minimize the risk of [Legionnaires' disease](#) and other diseases associated with water, [take steps](#) to ensure that all water systems and features (e.g., drinking fountains, decorative fountains) are safe to use after a prolonged facility shutdown. Drinking fountains should be cleaned and disinfected but encourage staff and players to bring their own water to minimize touching water fountains.
- **Modified Layouts and Social (Physical) Distancing**
  - Identify adult staff members or volunteers to help maintain [social distancing](#) among youth, coaches, umpires/referees, and spectators (if state and local directives allow for spectators).
  - Space players at least 6 feet apart on the field while participating in the sport (e.g., during warmup, skill building activities, simulation drills)
  - Discourage unnecessary physical contact, such as high fives, handshakes, fist bumps, or hugs.
  - Prioritize outdoor, as opposed to indoor, practice and play as much as possible.
  - Create distance between players when explaining drills or the rules of the game.
  - If keeping physical distance is difficult with players in competition or group practice, consider relying on individual skill work and drills.
  - Encourage players to wait in their cars with guardians until just before the beginning of a practice, warm-up, or game, instead of forming a group.
  - Limit the use of carpools or van pools. When riding in an automobile to a sports event, encourage players to ride to the sports event with persons living in their same household.
  - If practices or competition facilities must be shared, consider increasing the amount of time between practices and competitions to allow for one group to leave before another group enters the facility. If possible, allow time for cleaning and/or disinfecting.
- **Physical Barriers and Guides**
  - Provide physical guides, such as signs and tape on floors or playing fields, to make sure that coaches and players remain at least 6 feet apart.

# Dear Parents

- **Communal Spaces**

- Close shared spaces such as locker rooms, if possible; otherwise, stagger use and [clean and disinfect](#) between use.
- Limit the number of players sitting in confined player seating areas (e.g., dugouts) by allowing players to spread out into spectator areas if more space is available (e.g., if spectators are not allowed).

Youth sports organizations may consider implementing several strategies to maintain healthy operations.

- **Protections for Staff and Players at Higher Risk for Severe Illness from COVID-19**

- Offer options for individuals at [higher risk of severe illness from COVID-19](#) that limit exposure risk (such as virtual coaching and in-home drills).
- Limit youth sports participation to staff and youth who live in the local geographic area (e.g., community, city, town, or county) to reduce risk of spread from areas with higher levels of COVID-19.

- **Regulatory Awareness**

- Be aware of state or local regulatory agency policies related to group gatherings to determine if events can be held.

- **Identifying Small Groups and Keeping them Together (Cohorting)**

- Keep players together in small groups with dedicated coaches or staff, and make sure that each group of players and coach avoid mixing with other groups as much as possible. Teams might consider having the same group of players stay with the same coach or having the same group of players rotate among coaches.
- Consider staging within-team scrimmages instead of playing games with other teams to minimize exposure among players and teams.

- **Staggered Scheduling**

- Stagger arrival and drop-off times or locations by cohort (group) or put in place other protocols to limit contact between groups and with guardians as much as possible. One example is increasing the amount of time between practices and competitions to allow for one group to depart before another group enters the facility. This also allows for more time to clean the facility between uses.
- When possible, use flexible worksites (e.g., telework) and flexible work hours (e.g., staggered shifts) to help establish policies and practices for social distancing (maintaining a distance of approximately 6 feet) between employees and others, especially if social distancing is recommended by state and local health authorities.

- **Gatherings, Spectators, and Travel**

- Avoid group events, such as games, competitions, or social gatherings, where spacing of at least 6 feet between people cannot be maintained.
- Limit any nonessential visitors, spectators, volunteers, and activities involving external groups or organizations as much as possible – especially with individuals not from the local geographic area (e.g., community, town, city, or county).
- Avoid activities and events such as off-site competitions or excursions (e.g., watching a professional team compete).

# Dear Parents

- **Designated COVID-19 Point of Contact**
  - Designate a youth sports program staff person to be responsible for responding to COVID-19 concerns. All coaches, staff, officials, and families should know who this person is and how to contact them.
- **Communication Systems**
  - Put systems in place for:
    - Consistent with applicable law and privacy policies, having coaches, staff, umpires/officials, and families of players (as feasible) self-report to the youth sports organization if they have [symptoms](#) of COVID-19, a positive test for COVID-19, or were exposed to someone with COVID-19 within the last 14 days in accordance with [health information sharing regulations for COVID-19](#) (e.g. see “Notify Health Officials and Close Contacts” in the **Preparing for When Someone Gets Sick section below**), and other applicable laws and regulations.
    - Notifying staff, officials, families, and the public of youth sports facility closures and restrictions in place to limit COVID-19 exposure (e.g., limited hours of operation).
  - Develop policies for return-to-play after COVID-19 illness. CDC’s criteria to [discontinue home isolation and quarantine](#) can inform these policies.
- **Back-up Staffing Plan**
  - Monitor absenteeism of coaches and officials, cross-train staff, and create a roster of trained back-up personnel.
- **Coach and Staff Training**
  - Train coaches, officials, and staff on all safety protocols.
  - Conduct training virtually, or ensure that [social distancing](#) is maintained during training.
- **Recognize Signs and Symptoms**
  - If feasible, conduct daily health checks (e.g., [symptom checking](#)) of coaches, officials, staff, and players safely and respectfully, and in accordance with any applicable privacy and confidentiality laws and regulations.
  - Youth sports program administrators may use examples of screening methods found in CDC’s supplemental [Guidance for Child Care Programs that Remain Open](#) as a guide for screening children, and CDC’s [General Business FAQs](#) for screening staff.
- **Sharing Facilities**
  - Encourage any organizations that share or use the youth sports facilities to also follow these considerations.
- **Support Coping and Resilience**
  - Encourage employees to take breaks from watching, reading, or listening to news stories, including social media if they are feeling overwhelmed or distressed.
  - Promote healthy eating, exercising, getting sleep, and finding time to unwind.
  - Encourage employees to talk with people they trust about their concerns and how they are feeling.
  - Consider posting signs for the national distress hotline: 1-800-985-5990, or text Talk With Us to 66746

# Dear Parents

## Preparing for When Someone Gets Sick

Youth sports organizations may consider implementing several strategies to prepare for when someone gets sick.

- **Advise Sick Individuals of Home Isolation Criteria**
  - Sick coaches, staff members, umpires/officials, or players should not return until they have met CDC's [criteria to discontinue home isolation](#).
- **Isolate and Transport Those Who are Sick**
  - Make sure that coaches, staff, officials, players, and families know that sick individuals should not attend the youth sports activity, and that they should notify youth sports officials (e.g., the COVID-19 point of contact) if they (staff) or their child (families) become sick with COVID-19 [symptoms](#), test positive for COVID-19, or have been [exposed](#) to someone with COVID-19 symptoms or a confirmed or suspected case.
  - Immediately separate coaches, staff, officials, and players with COVID-19 [symptoms](#) (i.e., fever, cough, shortness of breath) at any youth sports activity. Individuals who are sick should go home or to a healthcare facility, depending on how severe their symptoms are, and follow [CDC guidance for caring for oneself and others](#) who are sick. Individuals who have had [close contact](#) with a person who has [symptoms](#) should be separated and sent home as well, and follow [CDC guidance for community-related exposure](#) (see “Notify Health Officials and Close Contacts” below). If symptoms develop, individuals and families should follow [CDC guidance for caring for oneself and others](#) who are sick.
  - Establish procedures for safely transporting anyone who is sick to their home or to a healthcare facility. If you are calling an ambulance or bringing someone to the hospital, try to call first to alert them that the person may have COVID-19.
- **Clean and Disinfect**
  - Close off areas used by a sick person and do not use these areas until after [cleaning and disinfecting](#) them (for outdoor areas, this includes surfaces or shared objects in the area, if applicable).
  - Wait at least 24 hours before cleaning and disinfecting. If 24 hours is not feasible, wait as long as possible. Ensure [safe and correct use](#) and storage of [cleaning external icon](#) and disinfection products, including storing them securely away from children.
- **Notify Health Officials and Close Contacts**
  - In accordance with state and local privacy and confidentiality laws and regulations, youth sports organizations should notify [local health officials](#), youth sports program staff, umpires/officials, and families immediately of any case of COVID-19 while maintaining confidentiality in accordance with the [Americans with Disabilities Act \(ADA\)](#) and other applicable laws and regulations.
  - Work with [local health officials](#) to develop a reporting system (e.g., letter) youth sports organizations can use to notify health officials and close contacts of cases of COVID-19.
  - Advise those who have had [close contact](#) with a person diagnosed with COVID-19 to stay home and [self-monitor for symptoms](#), and to follow [CDC guidance](#) if symptoms develop.

# Dear Parents

- Communication Resources Organized youth or adult recreational sports leagues in groups of up to 25 players, excluding coaches and referees/umpires, are permitted.
- Spectators are strongly discouraged for adult sports. Spectators, like parents, are permitted for youth sports, so long as members from different households maintain at least 6 feet of physical distance from each other.
- Organizers must maintain contact information and team rosters, and be prepared to support local public health contact tracing efforts if exposures occur.
- Consider participating in low or no-contact sports (like baseball, cross country, or cycling) instead of high-contact sports (like wrestling or soccer).
- Do not share snacks or water, except in emergency situations.
- Consider only holding games with other teams every two weeks, to minimize the number of new teams of players interacting. Regular practices with the same group are fine.
- Games that require extensive travel are strongly discouraged.
- Practice social distancing of 6 feet from other households during drop off/pick up of players.
- Masks are encouraged where feasible (i.e. in dugouts, by coaches, by spectators).
- Competitive events such as races and endurance events are allowed as long as social distancing and limitations on group size can be maintained. This includes implementing staggered start times and making efforts to prevent gatherings at starts and finishes, so that no more than 10 people are gathered at a time.
- Limit indoor facilities to up to 25% capacity, or 10 people, whichever is fewer, per room, so long as people can stay 6 feet apart from each other.
- Limit the pool to 25% capacity, or up to 10 people. It is strongly encouraged to limit pool activity to lap swim. If you permit open swim, make efforts to reduce in-pool interactions between people not in the same household.
- Teams participating in organized recreation against one another, and as a result, experiencing greater contact, should still observe the limit of 10 players, excluding coaches.
- Use a reservation system or use preexisting electronic capacity monitoring systems if feasible to space out and limit participants gathered at one time.
- Discourage use of any shared equipment and ensure all equipment is cleaned and disinfected between each use.
- Take steps, such as closing off a series of lockers to promote physical distancing and reduce gatherings in restrooms, showers and locker rooms.
- Provide access to hand sanitizer.
- Request staff members and patrons wear face coverings when they can do so safely.
- Conduct symptom and temperature checks for employees and refer symptomatic employees to the CDPHE Symptom Tracker.
- Employees who exhibit COVID-19 symptoms should not come to work.
- Employees who develop COVID-19 symptoms while at work should immediately notify their supervisor and be separated from others, sent home and referred to state or company support services.
- Post signs for employees and customers outlining good hand/respiratory hygiene and safety measures being taken